

GEOMETRIC REALIZATION AND K-THEORETIC DECOMPOSITION OF C*-ALGEBRAS

C. L. SCHOCHET

Mathematics Department
Wayne State University
Detroit, MI 48202

Submitted November 25, 1999

ABSTRACT. Suppose that A is a separable C^* -algebra and that G_* is a (graded) subgroup of the $\mathbb{Z}/2$ -graded group $K_*(A)$. Then there is a natural short exact sequence

$$(*) \quad 0 \rightarrow G_* \longrightarrow K_*(A) \longrightarrow K_*(A)/G_* \rightarrow 0.$$

In this note we demonstrate how to geometrically realize this sequence at the level of C^* -algebras. As a result, we KK -theoretically decompose A as

$$0 \rightarrow A \otimes \mathcal{K} \longrightarrow A_f \longrightarrow SA_t \rightarrow 0$$

where $K_*(A_t)$ is the torsion subgroup of $K_*(A)$ and $K_*(A_f)$ is its torsionfree quotient. Then we further decompose A_t : it is KK -equivalent to $\oplus_p A_p$ where $K_*(A_p)$ is the p -primary subgroup of the torsion subgroup of $K_*(A)$. We then apply this realization to study the Kasparov group $K^*(A)$ and related objects.

In Section 1 we produce the basic geometric realization. For any separable C^* -algebra A and group G_* we produce associated C^* -algebras A_s (s for subgroup) and A_q (q for quotient group) and, most importantly, a short exact sequence of C^* -algebras

$$0 \rightarrow A \otimes \mathcal{K} \rightarrow A_q \rightarrow SA_s \rightarrow 0$$

whose associated K_* -long exact sequence is $(*)$. In the case where G_* is the torsion subgroup of $K_*(A)$ we use the notation A_t (t for torsion) and A_f (f for torsionfree) respectively. We further decompose A_t into its p -primary summands A_p for each prime p .

Section 2 deals with the following question: may calculations of the Kasparov groups $KK_*(A, B)$ be reduced down to the four cases (A_t, B_t) , (A_t, B_f) , (A_f, B_t) and (A_f, B_f) ? We show that this is indeed possible in a wide variety of situations. Sections 3 and 4 deal with these special cases.

1991 *Mathematics Subject Classification.* Primary 46L80, 19K35, 46L85.

Key words and phrases. K -theory for C^* -algebras, geometric realization, Kasparov theory.

Geometric realization as a general technique was introduced to topological K -theory of spaces by M. F. Atiyah [1] in his proof of the Künneth theorem for $K^*(X \times Y)$. We adapted the technique [6] to prove the corresponding theorem for the K -theory for C^* -algebras and used it with J. Rosenberg in our proof of the Universal Coefficient Theorem (UCT) [4].

1. Geometric Realization

In this section we produce the main geometric realization and we extend the result to give a p -primary decomposition for a C^* -algebra.

Let $\mathcal{N}$ denote the bootstrap category [6, 4].

Theorem 1.1. *Suppose that A is a separable C^* -algebra. Let G_* be some subgroup of $K_*(A)$. Then there is an associated C^* -algebra $A_s \in \mathcal{N}$, a separable C^* -algebra A_q , and a short exact sequence*

$$(1.2) \quad 0 \rightarrow A \otimes \mathcal{K} \rightarrow A_q \rightarrow SA_s \rightarrow 0$$

whose induced K -theory long exact sequence fits into the commuting diagram

$$(1.3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & K_*(A_s) & \longrightarrow & K_*(A \otimes \mathcal{K}) & \longrightarrow & K_*(A_q) \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ 0 & \longrightarrow & G_* & \longrightarrow & K_*(A) & \longrightarrow & K_*(A)/G_* \longrightarrow 0. \end{array}$$

If A is nuclear then so is A_q . If $A \in \mathcal{N}$ then so is A_q . If $A \in \mathcal{N}$ and if G_* is a direct summand of $K_*(A)$ then A is KK -equivalent to $A_s \oplus A_q$.

Note that we think of A_s as realizing the subgroup G_* and A_q as realizing the quotient group $K_*(A)/G_*$, hence the notation.

Proof. Let A_s denote any C^* -algebra in $\mathcal{N}$ with

$$K_*(A_s) \cong G_*.$$

Such C^* -algebras exist and are unique up to KK -equivalence by the UCT [4]. Let

$$\theta : K_*(A_s) \rightarrow K_*(A)$$

be the corresponding homomorphism. Since $A_s \in \mathcal{N}$, the UCT holds for the pair (A_s, A) , and so θ is in the image of the index map

$$\gamma : KK_*(A_s, A) \rightarrow \text{Hom}_{\mathbb{Z}}(K_*(A_s), K_*(A)).$$

Say that

$$\theta = \gamma(\tau)$$

for some

$$\tau \in KK_0(A_s, A).$$

As A_s is nuclear,

$$KK_*(A_s, A) \simeq \text{Ext}(K_*(A_s), K_*(A))$$

and hence τ corresponds to an equivalence class of extensions of C^* -algebras of the form

$$0 \rightarrow A \otimes \mathcal{K} \rightarrow E \rightarrow SA_s \rightarrow 0.$$

Define $A_q = E$. (This choice depends upon the choice of A_s among its KK -equivalence class and the choice of τ modulo the kernel of γ). Note that E is nuclear/bootstrap if and only if A is nuclear/bootstrap. Then the diagram

$$\begin{array}{ccccccc} K_j(A_q) & \longrightarrow & K_j(SA_s) & \xrightarrow{\delta} & K_{j-1}(A \otimes \mathcal{K}) & \longrightarrow & K_{j-1}(A_q) \\ & & \downarrow \cong & & \downarrow \cong & & \\ & & K_{j-1}(A_s) & \xrightarrow{\theta} & K_{j-1}(A) & & \end{array}$$

commutes, and thus δ is mono and the long exact K_* -sequence breaks apart as shown.

If G_* is a direct summand of $K_*(A)$ then

$$K_*(A) \cong G_* \oplus K_*(A)/G_* \cong K_*(A_s) \oplus K_*(A_q) \cong K_*(A_s \oplus A_q)$$

and, replacing algebras by their suspensions as needed, the KK -equivalence is obtained.

□

Henceforth we shall regard A_s and A_q as C^* -algebras associated to A and G_* , with the understanding that these are well-defined only up to KK -equivalence modulo the kernel of γ , as explained above.

The next step is to decompose A_t into its p -primary components.

Theorem 1.4. *Let $A \in \mathcal{N}$ and suppose that $K_*(A)$ is a torsion group, so that $A = A_t$. Then A is KK -equivalent to a C^* -algebra $\oplus A_p$, where*

$$K_*(A_p) \cong K_*(A)_p$$

the p -primary torsion subgroup of $K_*(A)$.

Proof. For each prime p , choose $N_{(p)} \in \mathcal{N}$ with $K_1(N_{(p)}) = 0$ and

$$K_0(N_{(p)}) \cong \mathbb{Z}_{(p)}$$

the integers localized at p . Define

$$A_p = A_t \otimes N_{(p)}.$$

The Künneth formula [6] implies that

$$K_*(A_p) \cong K_*(A_t \otimes N_{(p)}) \cong K_*(A_t) \otimes K_*(N_{(p)}) \cong K_*(A_t) \otimes \mathbb{Z}_{(p)} \cong K_*(A)_p$$

as desired. Then

$$K_*(\oplus_p A_p) \cong \oplus_p K_*(A_p) \cong \oplus_p K_*(A)_p \cong K_*(A_t)$$

and another use of the UCT implies that A_t is KK -equivalent to $\oplus_p A_p$.

□

We summarize:

Theorem 1.5. *Suppose that A is a separable C^* -algebra. Then there is an associated C^* -algebra $A_t \in \mathcal{N}$, a separable C^* -algebra A_f , and a short exact sequence*

$$(1.6) \quad 0 \rightarrow A \otimes \mathcal{K} \rightarrow A_f \rightarrow SA_t \rightarrow 0$$

whose induced K -theory long exact sequence fits into the commuting diagram

$$(1.7) \quad \begin{array}{ccccccc} 0 & \longrightarrow & K_*(A_t) & \longrightarrow & K_*(A \otimes \mathcal{K}) & \longrightarrow & K_*(A_f) \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow \cong & & \downarrow \cong \\ 0 & \longrightarrow & K_*(A)_t & \longrightarrow & K_*(A \otimes \mathcal{K}) & \longrightarrow & K_*(A)_f \longrightarrow 0. \end{array}$$

If A is nuclear then so is A_f . If $A \in \mathcal{N}$ then so is A_f . Further, the C^* -algebra A_t has a p -primary decomposition: it is KK -equivalent to a C^* -algebra $\oplus_p A_p$, where $A_p \in \mathcal{N}$ for all p and

$$K_*(A_p) \cong K_*(A)_p$$

the p -primary torsion subgroup of $K_*(A)$. Finally, if $A \in \mathcal{N}$ and $K_*(A)_t$ is a direct summand of $K_*(A)$ then A may be replaced by the KK -equivalent C^* -algebra $A_t \oplus A_f$

□

2. Splitting the Kasparov Groups

If A and B are in $\mathcal{N}$ and their K -theory torsion subgroups $K_*(A)_t$ and $K_*(B)_t$ are direct summands then the final conclusion of Theorem 1.5 implies that we may reduce the computation of $KK_*(A, B)$ to the calculation of the four groups, namely

- (1) $KK_*(A_t, B_t)$
- (2) $KK_*(A_t, B_f)$
- (3) $KK_*(A_f, B_t)$
- (4) $KK_*(A_f, B_f)$.

We discuss the calculation of those groups in subsequent sections. In this section we see what can be done *without* assuming that the torsion subgroups are direct summands.

Theorem 2.1. *Suppose that $A \in \mathcal{N}$ and $K_*(B)$ is torsionfree. Then there is a short exact sequence*

$$(2.2) \quad 0 \rightarrow KK_*(A_f, B) \rightarrow KK_*(A, B) \rightarrow KK_*(A_t, B) \rightarrow 0.$$

In particular, letting $K^*(A) = KK_*(A, \mathbb{C})$ there is a short exact sequence

$$(2.3) \quad 0 \rightarrow K^*(A_f) \rightarrow K^*(A) \rightarrow K^*(A_t) \rightarrow 0.$$

If $K_*(B)$ is not necessarily torsionfree, then sequence 2.2 is exact if and only if the natural map

$$(2.4) \quad \theta_h^* : \text{Hom}_{\mathbb{Z}}(K_*(A), K_*(B)) \rightarrow \text{Hom}_{\mathbb{Z}}(K_*(A_t), K_*(B))$$

is onto, where $\theta : K_*(A_t) \rightarrow K_*(A)$ is the canonical inclusion.

Note that the map θ_h^* in (2.4) is frequently onto. This is the case, for instance, if $K_*(A_t)$ is a direct summand of $K_*(A)$.

The map θ is, up to isomorphism, the boundary homomorphism in the K_* -sequence associated to the short exact sequence

$$0 \rightarrow A \otimes \mathcal{K} \rightarrow A_f \rightarrow SA_t \rightarrow 0$$

and hence

$$\theta(x) = x \otimes_{A_t} \delta$$

where $\delta \in KK_1(A_t, A)$ by [9]. Thus the map θ_h^* of (2.4) is induced from a KK -pairing.

Proof. Consider the commuting diagram

$$\begin{array}{ccccccc}
 & \text{Hom}_{\mathbb{Z}}(K_*(A_t), K_*(B)) & & KK_{*-1}(A_t, B) & & 0 & \\
 & \beta \downarrow & & \downarrow & & \downarrow & \\
 0 \longrightarrow & \text{Ext}_{\mathbb{Z}}^1(K_*(A_f), K_*(B)) & \longrightarrow & KK_*(A_f, B) & \longrightarrow & \text{Hom}_{\mathbb{Z}}(K_*(A_f), K_*(B)) & \longrightarrow 0 \\
 & \downarrow & & \downarrow & & \downarrow & \\
 0 \longrightarrow & \text{Ext}_{\mathbb{Z}}^1(K_*(A), K_*(B)) & \longrightarrow & KK_*(A, B) & \longrightarrow & \text{Hom}_{\mathbb{Z}}(K_*(A), K_*(B)) & \longrightarrow 0 \\
 & \theta_e^* \downarrow & & \theta^* \downarrow & & \theta_h^* \downarrow & \\
 0 \longrightarrow & \text{Ext}_{\mathbb{Z}}^1(K_*(A_t), K_*(B)) & \longrightarrow & KK_*(A_t, B) & \longrightarrow & \text{Hom}_{\mathbb{Z}}(K_*(A_t), K_*(B)) & \longrightarrow 0 \\
 & \downarrow & & \downarrow & & \beta \downarrow & \\
 & 0 & & KK_{*+1}(A_f, B) & & \text{Ext}_{\mathbb{Z}}^1(K_*(A_f), K_*(B)) &
 \end{array}$$

The three middle rows are exact by the UCT, the middle column is exact by the exactness of KK , and the two outer columns are exact by the standard Hom-Ext -sequence.

Suppose that $K_*(B)$ is torsionfree. Then

$$(2.5) \quad \text{Hom}_{\mathbb{Z}}(K_*(A_t), K_*(B)) = 0$$

since $K_*(A_t)$ is a torsion group, and the surjectivity of θ_e^* implies the surjectivity of θ^* .

If $K_*(B)$ is not necessarily torsionfree, then the Snake Lemma [11] implies that there is an exact sequence

$$0 = \text{Coker}(\theta_e^*) \longrightarrow \text{Coker}(\theta^*) \longrightarrow \text{Coker}(\theta_h^*) \rightarrow 0$$

and hence θ_h^* is onto if and only if θ^* is onto. The theorem then follows immediately, for the middle column of the diagram degenerates to (2.2) if and only if θ^* is onto. $\square$

Next we consider the dual situation, when $K_*(A)$ is a torsion group.

Theorem 2.6. *Suppose that $A \in \mathcal{N}$ and that $K_*(A)$ is a torsion group. Then there is a natural exact sequence*

$$(2.7) \quad 0 \rightarrow KK_*(A, B_t) \rightarrow KK_*(A, B) \rightarrow KK_*(A, B_f) \rightarrow 0.$$

If $K_(A)$ is not a torsion group then sequence (*) is exact if and only if the natural map*

$$\pi_* : \text{Hom}_{\mathbb{Z}}(K_*(A), K_*(B)) \rightarrow \text{Hom}_{\mathbb{Z}}(K_*(A), K_*(B_f))$$

is onto, where $\pi : B \otimes \mathcal{K} \rightarrow B_f$ is the natural map.

The proof of this result is dual to that of Theorem 2.1 and is omitted for brevity. $\square$

3. Computing $KK_*(A_f, B)$

In this section we consider the case where $K_*(A)$ is torsionfree (so that $A = A_f$). Recall [2, 12] that a subgroup H of an abelian group K is *pure* if for each positive integer n ,

$$nH = H \cap nG,$$

and an extension of groups

$$0 \rightarrow H \rightarrow K \rightarrow G \rightarrow 0$$

is *pure* if H is a pure subgroup of K . For abelian groups G and H , $Pext_{\mathbb{Z}}^1(G, H)$ is the subgroup of $Ext_{\mathbb{Z}}^1(G, H)$ consisting of pure extensions.

Recall [5, 8] that there is a natural topology on the Kasparov groups and that with respect to this topology the UCT sequence splittings constructed in [4] are continuous, so that the splitting is a splitting of topological groups [9].

Theorem 3.1. *Suppose that $A \in \mathcal{N}$ and that $K_*(A)$ is torsionfree. Then there is a natural sequence of topological groups*

$$0 \rightarrow Pext_{\mathbb{Z}}^1(K_*(A), K_*(B)) \rightarrow KK_*(A, B) \rightarrow \text{Hom}_{\mathbb{Z}}(K_*(A), K_*(B)) \rightarrow 0$$

The group $Pext_{\mathbb{Z}}^1(K_(A), K_*(B))$ is the closure of zero in the natural topology on the group $KK_*(A, B)$ and thus the group $\text{Hom}_{\mathbb{Z}}(K_*(A), K_*(B))$ is the Hausdorff quotient of $KK_*(A, B)$.*

Proof. The UCT gives us the sequence

$$0 \rightarrow Ext_{\mathbb{Z}}^1(K_*(A), K_*(B)) \rightarrow KK_*(A, B) \rightarrow \text{Hom}_{\mathbb{Z}}(K_*(A), K_*(B)) \rightarrow 0$$

which splits unnaturally. If $K_*(A)$ is torsionfree then

$$Pext_{\mathbb{Z}}^1(K_*(A), K_*(B)) \cong Ext_{\mathbb{Z}}^1(K_*(A), K_*(B)).$$

The remaining part of the theorem holds since we have shown in general [10] that the group $Pext_{\mathbb{Z}}^1(K_*(A), K_*(B))$ is the closure of zero in the natural topology on $KK_*(A, B)$ in the presence of the UCT.

$\square$

We note that the resulting algebraic problems are frequently very difficult. If G is a torsionfree abelian group then $\text{Hom}_{\mathbb{Z}}(G, H)$ is unknown in general, though there is much known in special cases (cf. [2, 3]). The group $Pext_{\mathbb{Z}}^1(G, H)$ is also difficult, though the case $Pext_{\mathbb{Z}}^1(G, \mathbb{Z})$ is known (cf. [3]). We discuss $Pext$ in detail in [12].

4. Computing $KK_*(A_t, B)$

In this section we concentrate upon the situation when $K_*(A)$ is a torsion group. Before beginning, we digress slightly to recall [7] in more detail how one introduces coefficients into K -theory.

Given a countable abelian group G , select some C^* -algebra $N_G \in \mathcal{N}$ with

$$K_0(N_G) = G \quad K_1(N_G) = 0.$$

The C^* -algebra N_G is unique up to KK -equivalence, by the UCT. Then for any C^* -algebra A , define

$$(4.1) \quad K_j(A; G) \cong K_j(A \otimes N_G).$$

The Künneth Theorem [6] implies that there is a natural short exact sequence

$$(4.2) \quad 0 \rightarrow K_j(A) \otimes G \xrightarrow{\alpha} K_j(A; G) \rightarrow \text{Tor}_1^{\mathbb{Z}}(K_{j-1}(A), G) \rightarrow 0$$

which splits unnaturally. If G is torsionfree then α is an isomorphism

$$\alpha : K_j(A) \otimes G \xrightarrow{\cong} K_j(A; G).$$

Let $\mathbf{X}(G) = \text{Hom}(G, \mathbb{R}/\mathbb{Z})$ denote the Pontryagin dual of the group G .

Theorem 4.3. *Suppose that $A \in \mathcal{N}$ with $K_*(A)$ a torsion group and suppose that $K_*(B)$ is torsionfree, so that $A = A_t$ and $B = B_f$. Then:*

(1)

$$KK_*(A, B) \cong \text{Ext}_{\mathbb{Z}}^1(K_*(A), K_{*-1}(B)).$$

(2)

$$KK_*(A, B) \cong \text{Hom}_{\mathbb{Z}}(K_*(A), K_{*-1}(B) \otimes \mathbb{Q}/\mathbb{Z}).$$

(3) *The group $KK_*(A, B)$ is reduced and algebraically compact.*

(4)

$$K^j(A) \cong \mathbf{X}(K_{j-1}(A))$$

.

(5) *More generally, if $K_*(B)$ is finitely generated free, then*

$$KK_j(A, B) \cong \oplus_n \mathbf{X}(K_{j-1}(A))$$

where n is the number of generators of $K_(B)$.*

Proof. Part 1) follows at once from the UCT and the fact that there are no non-trivial homomorphisms from a torsion group to a torsionfree group. Part 2) follows from Part 1) by elementary homological algebra. Part 3) follows easily from a deep result of Fuchs and Harrison [cf. 2, 46.1]: if G is a torsion group then any group of the form $\text{Hom}_{\mathbb{Z}}(G, H)$ is reduced and algebraically compact. Part 4) follows from part 3) by setting $B = \mathbb{C}$ and observing that for any torsion group G , we have

$$\mathbf{X}(G) = \text{Hom}_{\mathbb{Z}}(G, \mathbb{Q}/\mathbb{Z}).$$

□

There is one additional case that fits into the present discussion and which partially overlaps with the result above.

Theorem 4.4. *Suppose that $A \in \mathcal{N}$ and that $K_*(A)$ has no free direct summand. Then there is a natural short exact sequence of topological groups*

$$(4.5) \quad 0 \rightarrow \operatorname{Hom}_{\mathbb{Z}}(K_*(A), \mathbb{R}) \rightarrow \mathbf{X}(K_*(A)) \xrightarrow{\chi} K^*(A) \rightarrow 0.$$

The map $\chi : \mathbf{X}(K_(A)) \rightarrow K^*(A)$ is a degree one continuous open surjection. It is a homeomorphism if and only if $K_*(A)$ is a torsion group.*

To be explicit about the grading,

$$\chi : \mathbf{X}(K_j(A)) \rightarrow K^{j-1}(A)$$

which is the usual parity shift as torsion phenomena move from homology to cohomology.

Proof. The UCT for $K^*(A)$ has the form

$$0 \rightarrow \operatorname{Ext}_{\mathbb{Z}}^1(K_*(A), \mathbb{Z}) \xrightarrow{\delta} K^*(A) \rightarrow \operatorname{Hom}_{\mathbb{Z}}(K_*(A), \mathbb{Z}) \rightarrow 0$$

with δ of degree one, so it suffices to compute Ext . In general the short exact sequence

$$0 \rightarrow \mathbb{Z} \rightarrow \mathbb{R} \rightarrow \mathbb{R}/\mathbb{Z} \rightarrow 0$$

yields a long exact sequence

$$\operatorname{Hom}_{\mathbb{Z}}(K_*(A), \mathbb{Z}) \rightarrow \operatorname{Hom}_{\mathbb{Z}}(K_*(A), \mathbb{R}) \rightarrow \mathbf{X}(K_*(A)) \rightarrow \operatorname{Ext}_{\mathbb{Z}}^1(K_*(A), \mathbb{Z}) \rightarrow 0.$$

The fact that $K_*(A)$ has no free direct summand implies that $\operatorname{Hom}_{\mathbb{Z}}(K_*(A), \mathbb{Z}) = 0$, so the sequence degenerates to

$$0 \rightarrow \operatorname{Hom}_{\mathbb{Z}}(K_*(A), \mathbb{R}) \rightarrow \mathbf{X}(K_*(A)) \rightarrow \operatorname{Ext}_{\mathbb{Z}}^1(K_*(A), \mathbb{Z}) \rightarrow 0.$$

Applying the UCT one obtains the sequence 4.5 as desired. The map χ is the composite of the UCT map and a natural homeomorphism. The rest of the Theorem is immediate.

□

REFERENCES

- [1] M. F. Atiyah, *Vector bundles and the Künneth formula*, Topology **1** (1962), 245-248.
- [2] László Fuchs, *Infinite Abelian Groups*, Pure and Applied Mathematics No. 36 , vol. 1, Academic Press, New York, 1970, pp. 290.
- [3] C. U. Jensen, *Les Foncteurs Dérivés de $\varprojlim$ et leur Applications en Théorie des Modules*, Lecture Notes in Mathematics , vol. 254, Springer, Verlag, New York, 1972.
- [4] J. Rosenberg and C. Schochet, *The Künneth theorem and the universal coefficient theorem for Kasparov's generalized K-functor*, Duke Math. J. **55** (1987), 431-474.
- [5] N. Salinas, *Relative quasidiagonality and KK-theory*, Houston J. Math. **18** (1992), 97-116.
- [6] C. Schochet, *Topological methods for C*-algebras II: geometric resolutions and the Künneth formula*, Pacific J. Math. **98** (1982), 443-458.
- [7] C. Schochet, *Topological methods for C*-algebras IV: mod p homology*, Pacific J. Math. **114** (1984), 447-468.
- [8] C. Schochet, *The fine structure of the Kasparov groups I: continuity of the KK-pairing*, submitted.
- [9] C. Schochet, *The fine structure of the Kasparov groups II: topologizing the UCT*, submitted.
- [10] C. Schochet, *The fine structure of the Kasparov groups III: relative quasidiagonality*, submitted.
- [11] C. Schochet, *The topological snake lemma and Corona algebras*, New York J. Math. **5** (1999), 131-137.
- [12] C. Schochet, *A Pext Primer: Pure extensions and $\lim^1$ for infinite abelian groups*, submitted.